

## WEST Search History

DATE: Monday, December 18, 2006

<b>Hide?</b>	<b><u>Set Name</u></b>	<b><u>Query</u></b>	<b><u>Hit Count</u></b>
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L4	L3 and desloratadine	49
<input type="checkbox"/>	L3	514/290	837
<input type="checkbox"/>	L2	L1 and desloratadine	7
<input type="checkbox"/>	L1	546/93	283

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1612RXD

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 AUG 09 INSPEC enhanced with 1898-1968 archive  
NEWS 4 AUG 28 ADISCTI Reloaded and Enhanced  
NEWS 5 AUG 30 CA(SM)/CAplus(SM) Austrian patent law changes  
NEWS 6 SEP 11 CA/CAplus enhanced with more pre-1907 records  
NEWS 7 SEP 21 CA/CAplus fields enhanced with simultaneous left and right  
truncation  
NEWS 8 SEP 25 CA(SM)/CAplus(SM) display of CA Lexicon enhanced  
NEWS 9 SEP 25 CAS REGISTRY(SM) no longer includes Concord 3D coordinates  
NEWS 10 SEP 25 CAS REGISTRY(SM) updated with amino acid codes for pyrrolysine  
NEWS 11 SEP 28 CEABA-VTB classification code fields reloaded with new  
classification scheme  
NEWS 12 OCT 19 LOGOFF HOLD duration extended to 120 minutes  
NEWS 13 OCT 19 E-mail format enhanced  
NEWS 14 OCT 23 Option to turn off MARPAT highlighting enhancements available  
NEWS 15 OCT 23 CAS Registry Number crossover limit increased to 300,000 in  
multiple databases  
NEWS 16 OCT 23 The Derwent World Patents Index suite of databases on STN  
has been enhanced and reloaded  
NEWS 17 OCT 30 CHEMLIST enhanced with new search and display field  
NEWS 18 NOV 03 JAPIO enhanced with IPC 8 features and functionality  
NEWS 19 NOV 10 CA/CAplus F-Term thesaurus enhanced  
NEWS 20 NOV 10 STN Express with Discover! free maintenance release Version  
8.01c now available  
NEWS 21 NOV 13 CA/CAplus pre-1967 chemical substance index entries enhanced  
with preparation role  
NEWS 22 NOV 20 CAS Registry Number crossover limit increased to 300,000 in  
additional databases  
NEWS 23 NOV 20 CA/CAplus to MARPAT accession number crossover limit increased  
to 50,000  
NEWS 24 DEC 01 CAS REGISTRY updated with new ambiguity codes  
NEWS 25 DEC 11 CAS REGISTRY chemical nomenclature enhanced  
NEWS 26 DEC 14 WPIDS/WPINDEX/WPIX manual codes updated  
NEWS 27 DEC 14 GBFULL and FRFULL enhanced with IPC 8 features and  
functionality  
  
NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items

NEWS IPC8      For general information regarding STN implementation of IPC 8  
NEWS X25      X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 11:33:29 ON 18 DEC 2006

=> file registry

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 11:33:42 ON 18 DEC 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 15 DEC 2006 HIGHEST RN 915749-75-6

DICTIONARY FILE UPDATES: 15 DEC 2006 HIGHEST RN 915749-75-6

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 30, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10510619.str

L1      STRUCTURE UPLOADED

=> s l1 ful

FULL SEARCH INITIATED 11:34:00 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED -      3543 TO ITERATE

100.0% PROCESSED      3543 ITERATIONS

826 ANSWERS

SEARCH TIME: 00.00.01

L2      826 SEA SSS FUL L1

10510619

=> file caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
166.94	167.15

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 11:34:03 ON 18 DEC 2006  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 18 Dec 2006 VOL 145 ISS 26  
FILE LAST UPDATED: 17 Dec 2006 (20061217/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s l2 and pure  
1287 L2  
440559 PURE  
17 PURES  
440568 PURE  
(PURE OR PURES)

L3 15 L2 AND PURE

=> s l3 and 99.9  
295294 99  
1878490 9  
19884 99.9  
(99(W)9)

L4 0 L3 AND 99.9

=> s l3 and 99  
295294 99

L5 1 L3 AND 99

=> d abs bib hitstr


L5 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN  
AB The present invention provides substantially pure desloratadine having an HPLC purity greater than 99.5% and having an absorbance less than 0.15 Au at 420 nm for a 5 % weight/volume solution in methanol, which does not show a peak for an impurity at a relative retention time in the range from 0.85 to 0.99 (relative to desloratadine appearing at a retention time of 25±5 min), which is greater than the discard limit set at less than 0.025 % of the total area,

when tested according to an HPLC method performed using a Hypersil BDS C8 column (15 cm x 4.6 mm, 5 µm particle size) with the following parameters: Mobile phase: a buffer solution having a pH of about 3, methanol and acetonitrile in a volume ratio of 8:1:1; Injection volume: 20 µL; Flow rate: 1.5 mL/min; Run time: 75 min; Discard limit: Set at less than 0.025% of total area. The present invention also provides a process for the preparation of substantially pure desloratadine by the process comprising acidic hydrolysis of desloratadine analogs with a strong organic acid or a mineral acid.

AN 2003:836755 CAPLUS  
 DN 139:328323  
 TI Substantially pure antihistaminic compound  
 IN Thennati, Rajamannar; Chitturi, Trinadha Rao; Kanangi, Shivramchandra; Unnam, Raja Sekhar; Jadav, Kanaksinh Jesingbhai  
 PA Sun Pharmaceutical Industries Limited, India  
 SO PCT Int. Appl., 19 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003086275	A2	20031023	WO 2003-IN156	20030416
	WO 2003086275	A3	20040212		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2003262141	A1	20031027	AU 2003-262141	20030416
	US 2006058334	A1	20060316	US 2004-510619	20041007
PRAI	IN 2002-MU348	A	20020415		
	WO 2003-IN156	W	20030416		


OS MARPAT 139:328323  
 IT 100643-71-8P, Desloratadine  
 RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (preparation of substantially pure desloratadine by acid hydrolysis)  
 RN 100643-71-8 CAPLUS  
 CN 5H-Benzo[5,6]cyclohepta[1,2-b]pyridine, 8-chloro-6,11-dihydro-11-(4-piperidinylidene)- (9CI) (CA INDEX NAME)



```

IT      79794-75-5
        RL: RCT (Reactant); RACT (Reactant or reagent)
            (preparation of substantially pure desloratadine by acid
            hydrolysis)
RN      79794-75-5  CAPLUS
CN      1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-
        benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI)  (CA
        INDEX NAME)

```



```
=> d l3 abs bib fhitrstr 1-15
```

L3 ANSWER 1 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
AB A review. Since the occurrence of allergic diseases increases, the need for antihistaminic drugs also grows. This requires new and more effective methods for their anal. Electromigration techniques are now often used in pharmaceutical anal. and can complement the use of HPLC. Data from the last 15 yr on the use of electromigration methods (capillary zonal electrophoresis, isotachopheresis, micellar electrokinetic chromatog., capillary electrochromatog.) in the separation and determination of

## H1-antihistaminics

in pure form, drugs, and biol. material are discussed. Data on the anal. of azelastine, cyclizine, diphenylhydramine, dioxopromethazine, dimetindene, embramine, pheniramine, hydroxyzine, meclozine, loratidine, and promethazine are examined

AN 2006:90923 CAPLUS

DN 145:174508

TI The use of electromigration methods in analysis of H1-antihistaminics

AU Kubacak, P.; Mikus, P.; Valaskova, I.; Havranek, E.

CS Farm. Fak., Univ. Komenského, Bratislava, Slovakia

SO Farmaceutický Obzor (2005), 74 (11), 301-306

CODEN: FAOBAS; ISSN: 0014-8172

PB Zdravotnicke Vydavatelstvo HERBA, spol. s r. o.

DT Journal; General Review


LA Slovak

IT 79794-75-5, Loratidine

RL: ANT (Analyte); ANST (Analytical study)  
(electromigration methods use in anal. of H1-antihistaminic drugs)

RN 79794-75-5 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)



L3 ANSWER 2 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB A rapid, sensitive, accurate, precise, reproducible, and versatile method for determining the purity of reference drug stds. and the routine anal. of illicit

drugs and adulterants using proton ( $^1\text{H}$ ) NMR Spectroscopy is presented.

The methodol. uses a weighed sample dissolved in a deuterated solvent or solvent mixture containing a high purity internal standard The NMR experiment

employs 8

scans using a 45 s delay and  $90^\circ$  pulse. In the determination of purity of reference stds., the number of quant. detns. available is equal to the number

of peak

groups that are baseline resolved. The relative standard deviation (RSD) of these signals is usually  $<1\%$  for pure stds., and the results

agree well with other purity determining methods. This method can also aid in the determination of correct mol. weight for stds. containing an unknown

number of waters of

hydration or an unknown number of acids per drug in salts. Because the molar response for the hydrogen nucleus is 1 for all compds., and since no separation media are used, only one linearity study is required to test a probe. In the presented study, the linearity of the NMR probe was determined using methamphetamine HCl dissolved in deuterium oxide ( $\text{D}_2\text{O}$ ) with maleic acid as the internal standard (5 mg) for a range of concns. from 0.033 to 69.18 mg/mL with a resulting correlation coefficient of  $>0.9999$  for all 6 methamphetamine peak groups. The spectra of complex illicit heroin, methamphetamine, MDMA, and cocaine samples are presented, as well as an extensive list of compds., their solubilities and the solvent(s) and internal standard used.

AN 2005:1277701 CAPLUS

DN 144:227583

TI Proton nuclear magnetic resonance spectroscopy (NMR) methods for determining the purity of reference drug standards and illicit forensic drug seizures

AU Hays, Patrick A.

CS Drug Enforcement Administration, Special Testing and Research Laboratory, U.S. Department of Justice, Dulles, VA, 20166, USA

SO Journal of Forensic Sciences (2005), 50(6), 1342-1360  
CODEN: JFSCAS; ISSN: 0022-1198

PB ASTM International

DT Journal

LA English


IT 79794-75-5, Loratadine

RL: ANT (Analyte); ANST (Analytical study)

and (proton NMR spectroscopy for purity determination of reference drug stds.  
illicit forensic drug seizures)

RN 79794-75-5 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)



RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 3 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB The inhibitory effects of different antihistamines on the expression of chemokine of cultivated skin in vitro was investigated. The skin was cultivated with mizolastine, cetirizine, loratadine, fexofenadine, dexamethasone and cyclosporin A, resp. Meanwhile histamine and arachidonic acid were added. Twenty-four hours later the total RNAs were obtained and purified with tri-pure isolation reagent. The mRNA expression levels of monocyte chemotactic protein-1 (MCP-1), monocyte chemotactic protein-3 (MCP-3), regulated on activation, normal T cell expressed, and secreted (RANTES) and eotaxin were analyzed by RT-PCR methods. The concns. of interleukin (IL)-4, IL-5 and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) in cultivated supernatant were detected by ELISA. In the mizolastine group the MCP-1 mRNA expression was significantly lower than that in cetirizine, loratadine or fexofenadine group and the MCP-3 mRNA expression was significantly lower than that in cetirizine or loratadine group ( $P < 0.05$ ). The RANTES mRNA expression and the eotaxin mRNA expression of the mizolastine group was significantly lower than that in loratadine group ( $P < 0.05$ ). The level of IL-4 in supernatant in mizolastine group was significantly lower than in fexofenadine group, and the level of IL-5 was significantly lower than that in cetirizine, loratadine and fexofenadine groups, while the level of TNF- $\alpha$  was significantly lower than that in loratadine group. Mizolastine inhibited the expression of chemokines and cytokines more strongly than the other antihistamines.

AN 2005:1140035 CAPLUS

DN 143:415811

TI Inhibition effects of antihistamines for chemokines and cytokines expression of cultivated skin in vitro


AU Pan, Hu; Hao, Fei

CS Southwest Hospital, Third Military Medical University, Chongqing, 400038, Peop. Rep. China


SO Linchuang Pifuke Zazhi (2005), 34(1), 20-22



CODEN: LPZAEH; ISSN: 1000-4963  
PB Linchuang Pifuke Zazhi Bianjibu  
DT Journal  
LA Chinese  
IT 79794-75-5, Loratadine  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
(Biological study); USES (Uses)  
(inhibition effects of antihistamines for chemokines and cytokines  
expression of cultivated skin in vitro)  
RN 79794-75-5 CAPLUS  
CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-  
benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA  
INDEX NAME)



L3 ANSWER 4 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
AB Two simple extractive spectrophotometric methods were developed for the  
estimation of loratadine in pure and pharmaceutical dosage forms.  
These methods are based on the formation of complexes of the drug with the  
2 acid dyes Orange-II and Eriochrome Black-T in acidic medium followed by  
their extraction into chloroform. The absorbance of the chloroform exts. is  
measured against the corresponding reagent blanks. These methods were  
statistically evaluated and found to be precise and accurate.  
AN 2005:261705 CAPLUS  
DN 142:322948  
TI Assay of loratadine by extractive spectrophotometry  
AU Rao, J. V. L. N. Seshagiri; Rao, Y. Srinivasa; Murthy, T. K.;  
Jitendrababu, V.; Chowdary, K. P. R.  
CS Department Of Pharmaceutical Sciences, Andhra University, Visakhapatnam,  
530 003, India  
SO Acta Ciencia Indica, Chemistry (2003), 29(3), 207-209  
CODEN: ACICDV; ISSN: 0253-7338  
PB Pragati Prakashan  
DT Journal  
LA English  
IT 79794-75-5, Loratadine  
RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL  
(Biological study); USES (Uses)  
(assay of loratadine by extractive spectrophotometry)  
RN 79794-75-5 CAPLUS  
CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-  
benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA  
INDEX NAME)




RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 5 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB A review. Treatment for allergic conjunctivitis has markedly expanded in recent years, providing opportunities for more focused therapy, but often leaving both physicians and patients confused over the variety of options. As monotherapy, oral antihistamines are an excellent choice when attempting to control multiple early-phase, and some late-phase, allergic symptoms in the eyes, nose and pharynx. Unfortunately, despite their efficacy in relief of allergic symptoms, systemic antihistamines may result in unwanted adverse effects, such as drowsiness and dry mouth. Newer second-generation antihistamines (cetirizine, fexofenadine, loratadine and desloratadine) are preferred over older first-generation antihistamines in order to avoid the sedative and anticholinergic effects that are associated with first-generation agents. When the allergic symptom or complaint, such as ocular pruritus, is isolated, focused therapy with topical (ophthalmic) antihistamines. Is often efficacious and clearly superior to systemic antihistamines, either as monotherapy or in conjunction with an oral or intranasal agent. Topical antihistaminic agents not only provide faster and superior relief than systemic antihistamines, but they may also possess a longer duration of action than other classes including vasoconstrictors, pure mast cell stabilizers, NSAIDs and corticosteroids. Many antihistamines have anti-inflammatory properties as well. Some of this anti-inflammatory effect seen with 'pure' antihistamines (levocabastine and emedastine) may be directly attributed to the blocking of the histamine receptor that has been shown to downregulate intercellular adhesion mol.-1 expression and, in turn, limit chemotaxis of inflammatory cells. Some topical multiple-action histamine H1-receptor antagonists (olopatadine, ketotifen, azelastine and epinastine) have been shown to prevent activation of neutrophils, eosinophils and macrophages, or inhibit release of leukotrienes, platelet-activating factors and other inflammatory mediators. Topical vasoconstrictor agents provide rapid relief, especially for redness; however, the relief is often short-lived, and overuse of vasoconstrictors may lead to rebound hyperemia and irritation. Another class of topical agents, mast cell stabilizers (sodium cromoglycate [cromolyn sodium], nedocromil and lodoxamide), may be considered; however, they generally have a much slower onset of action. The efficacy of mast cell stabilizers may be attributed to anti-inflammatory properties in addition to mast cell stabilization. In the class of topical NSAIDs, ketorolac has been promoted for ocular itching but has been found to be

inferior for relief of allergic conjunctivitis when compared with olopatadine and emedastine. Lastly, topical corticosteroids may be considered for severe seasonal ocular allergy symptoms, although long-term use should be avoided because of risks of ocular adverse effects, including glaucoma and cataract formation.

AN 2005:153798 CAPLUS  
 DN 142:456071  
 TI Efficacy and tolerability of newer antihistamines in the treatment of allergic conjunctivitis  
 AU Bielory, Leonard; Lien, Kenneth W.; Bigelsen, Steve  
 CS Department of Medicine, Pediatrics and Ophthalmology, Division of Allergy, Immunology and Rheumatology, Immuno-Ophthalmology Service, UMDNJ-New Jersey Medical School, Newark, NJ, USA  
 SO Drugs (2005), 65(2), 215-228  
 CODEN: DRUGAY; ISSN: 0012-6667  
 PB Adis International Ltd.  
 DT Journal; General Review  
 LA English  
 IT 79794-75-5, Loratadine  
 RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (antihistamines in treatment of allergic conjunctivitis)  
 RN 79794-75-5 CAPLUS  
 CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)



RE.CNT 110 THERE ARE 110 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 6 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
 AB The invention relates to novel microemulsions comprising a non-polar lipid, at least one polar solvent, a surfactant, and a polar lipid. A microemulsion of these ingredients provides an environment that substantially encloses airborne particles, i.e., allergens, bacteria and viruses, and it can be used for entrapping such particles. The inventive microemulsions are especially adapted for the prevention of symptoms in mammals, which are indirectly or directly caused by airborne particles. The invention further provides compns. comprising such microemulsions and methods of treatment comprising administering the same. For example, a microemulsion was prepared, containing glycerol monooleate 34%, propylene glycol

23%, polyethylene glycol 400 18%, sesame oil 11%, saline solution 10%, and polysorbate 80 4%. The microemulsion was administered as a spray to human volunteers. No signs or complaints of adverse effects were made. The formulation showed a superior wetting as compared with sesame oil and pure saline. Similar formulations protected MDCK and HSG cells in the culture from infection with influenza A virus.

AN 2005:55045 CAPLUS

DN 142:141249

TI Microemulsions comprising lipids for preventing airway diseases

IN Cockbain, Julian; Wollmer, Per; Andersson, Morgan; Greiff, Lennart; Landh, Thomas

PA Nares AB, Swed.

SO PCT Int. Appl., 36 pp.


CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005004843	A1	20050120	WO 2004-GB2925	20040707
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1648412	A1	20060426	EP 2004-743268	20040707
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
	US 2006269578	A1	20061130	US 2006-563828	20060508
PRAI	SE 2003-2005	A	20030707		
	SE 2003-2006	A	20030707		
	US 2003-484664P	P	20030707		
	WO 2004-GB2925	W	20040707		
IT	79794-75-5, Loratadine				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (microemulsions comprising lipids, polar solvent, and surfactant for entrapping airborne particles and preventing airway diseases)				
RN	79794-75-5 CAPLUS				
CN	1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)				



RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 7 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB A consumable film adapted to adhere to and dissolve in the oral cavity of a warm-blooded animal including humans, comprising a modified starch, pharmaceutically active agent and, optionally, at least one water-soluble polymer, such as pullulan, cellulose derivs., polyvinylpyrrolidone, acacia gum, poly(acrylic acid), etc. For example, preparation of a quick dissolving consumable film containing 15.0 mg dextromethorphan-HBr and Pure -Cote B793 base, using xanthan gum, locust bean gum, carrageenan and Pure-Cote B793 as film-forming ingredients was presented.

AN 2004:965039 CAPLUS

DN 141:400962

TI Fast dissolving orally consumable films containing a modified starch for improved heat and moisture resistance

IN Fadden, David John; Kulkarni, Neema Mahesh; Sorg, Albert Frank

PA Warner-Lambert Company LLC, USA

SO PCT Int. Appl., 33 pp.

CODEN: PIXXD2


DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2004096193	A1	20041111	WO 2004-IB1398	20040419
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:				
BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004233744	A1	20041111	AU 2004-233744	20040419
CA 2523372	A1	20041111	CA 2004-2523372	20040419
EP 1622595	A1	20060208	EP 2004-728218	20040419
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004009463	A	20060502	BR 2004-9463	20040419
CN 1784218	A	20060607	CN 2004-80011888	20040419

JP 2006525308 T 20061109 JP 2006-506579 20040419  
 US 2004247648 A1 20041209 US 2004-838045 20040503  
 PRAI US 2003-467339P P 20030502  
 WO 2004-IB1398 A 20040419  
 IT 79794-75-5, Loratadine  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (fast dissolving oral films containing modified starch and optionally  
 water-soluble polymer for improved heat and moisture resistance)  
 RN 79794-75-5 CAPLUS  
 CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-  
 benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA  
 INDEX NAME)




RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 8 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
 AB Disclosed is a free-flowing solid formulations of drugs or pharmaceutical  
 agents which have poor aqueous solubility are obtained by admixing a liquid or  
 gel  
 composition that includes 1-30 % of the drug, 5-60 % of a surfactant, 10-40 %  
 of water; 1-20 % of unsatd. fatty acid ester, 0-50 % water miscible  
 pharmaceutically acceptable polyol and 1-10 % phospholipid with a  
 pharmaceutically acceptable suitable solid carrier and thereafter drying  
 the admixt. The free-flowing powder is suitable for being formed into  
 tablets or capsules. The drug or pharmaceutical agent is solubilized in  
 the formulation and has significantly improved bio-availability when  
 compared to the drug tested in its pure form. A gel composition  
 containing polyoxyethylene sorbitan monooleate 35, propylene glycol 25, Et  
 linoleate 8, simvastatin 4, and 5 % lecithin aqueous solution q.s. to 100 % was  
 formulated. Colloidal silicon dioxide 30 parts was granulated with the  
 obtained gel 70 parts. The granules was dried to provide a free-flowing  
 powder. When this powder was exposed to a gastric medium of pH 1.2, 67 %  
 of the drug simvastatin dissolved within 10 min.  
 AN 2004:490267 CAPLUS  
 DN 141:42919  
 TI Free-flowing solid formulations with improved bio-availability of poorly  
 water soluble drugs and process for making the same  
 IN Li, Wenji; Alosio, Edward; Dema-Ala, Bricini Faith; Nguyen, Amy  
 PA USA  
 SO U.S. Pat. Appl. Publ., 9 pp.  
 CODEN: USXXCO  
 DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004115226	A1	20040617	US 2002-317657	20021212
	WO 2004054540	A2	20040701	WO 2003-US38979	20031209
	WO 2004054540	A3	20040930		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003300833	A1	20040709	AU 2003-300833	20031209
	JP 2006511536	T	20060406	JP 2004-560372	20031209
	US 2006263397	A1	20061123	US 2006-494131	20060727
PRAI	US 2002-317657	A	20021212		
	WO 2003-US38979	W	20031209		
IT	79794-75-5, Loratadine				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (free-flowing solid formulations with improved bio-availability of poorly water soluble drugs obtained from gel compns. containing surfactants, fatty acid esters, polyols, and phospholipids)				
RN	79794-75-5 CAPLUS				
CN	1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)				



L3 ANSWER 9 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
GI



AB Title compds. I [R1, R2, R3 = halo, alkyl, OR4; R4 = alkyl; X = H] were prepared via the reduction (+) piperidinylienes II. For example, diisobutylaluminum hydride reduction of atropisomer (+) II (R1, R3 = Br; R2 = Cl; X = H), e.g., prepared from piperidinecarboxylic acid II (R1, R3 = Br; R2 = Cl; X = CO2Et) in 2-steps, afforded claimed piperidine I (R1, R3 = Br; R2 = Cl; X = H) in  $[\alpha] = +51.9^\circ$ . Of note, atropisomer (-) II (R1, R3 = Br; R2 = Cl; X = H) was converted in 45% yield to (+) II (R1, R3 = Br; R2 = Cl; X = H) via heating in 1,2-dichlorobenzene at  $150^\circ\text{C}$  for 7-days. Compds. I are useful intermediates in the preparation of farnesyl protein transferase (FPT) inhibitor SCH66336.

AN 2004:212161 CAPLUS

DN 140:270738

TI Preparation of optically pure atropisomers of the piperidinylienebenzocycloheptapyridine intermediate of SCH66336

IN Njoroge, F. George; Vibulbhan, Bancha; Girijavallabhan, Viyyoor M.

PA Schering Corporation, USA

SO U.S., 7 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6706883	B1	20040316	US 1999-345966	19990701
PRAI	US 1998-91585P	P	19980702		
OS	CASREACT 140:270738; MARPAT 140:270738				
IT	218453-59-9P				


RL: PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)

(preparation of optically pure atropisomers of the piperidinylienebenzocycloheptapyridine intermediate of SCH66336)

RN 218453-59-9 CAPLUS

CN 5H-Benzo[5,6]cyclohepta[1,2-b]pyridine, 3,10-dibromo-8-chloro-6,11-dihydro-11-(4-piperidinyliene)-, (+)- (9CI) (CA INDEX NAME)





RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 10 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB The present invention provides substantially pure desloratadine having an HPLC purity greater than 99.5% and having an absorbance less than 0.15 Au at 420 nm for a 5 % weight/volume solution in methanol, which does not show a peak for an impurity at a relative retention time in the range from 0.85 to 0.99 (relative to desloratadine appearing at a retention time of 25±5 min), which is greater than the discard limit set at less than 0.025 % of the total area, when tested according to an HPLC method performed using a Hypersil BDS C8 column (15 cm x 4.6 mm, 5 µm particle size) with the following parameters: Mobile phase: a buffer solution having a pH of about 3, methanol and acetonitrile in a volume ratio of 8:1:1; Injection volume: 20 µL; Flow rate: 1.5 mL/min; Run time: 75 min; Discard limit: Set at less than 0.025% of total area. The present invention also provides a process for the preparation of substantially pure desloratadine by the process comprising acidic hydrolysis of desloratadine analogs with a strong organic acid or a mineral acid.

AN 2003:836755 CAPLUS

DN 139:328323

TI Substantially pure antihistaminic compound

IN Thennati, Rajamannar; Chitturi, Trinadha Rao; Kanangi, Shivramchandra; Unnam, Raja Sekhar; Jadav, Kanaksinh Jesingbhai

PA Sun Pharmaceutical Industries Limited, India

SO PCT Int. Appl., 19 pp.

CODEN: PIXXD2


DT Patent

LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003086275	A2	20031023	WO 2003-IN156	20030416
	WO 2003086275	A3	20040212		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2003262141	A1	20031027	AU 2003-262141	20030416

US 2006058334 A1 20060316 US 2004-510619 20041007  
 PRAI IN 2002-MU348 A 20020415  
 WO 2003-IN156 W 20030416  
 OS MARPAT 139:328323  
 IT 100643-71-8P, Desloratadine  
 RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use);  
 BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (preparation of substantially pure desloratadine by acid  
 hydrolysis)  
 RN 100643-71-8 CAPLUS  
 CN 5H-Benzo[5,6]cyclohepta[1,2-b]pyridine, 8-chloro-6,11-dihydro-11-(4-  
 piperidinylidene)- (9CI) (CA INDEX NAME)



L3 ANSWER 11 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
 AB Three simple and accurate methods are presented for determination of  
 Cetirizine,  
 Fexofenadine, Loratadine and Acrivastine in pure form and com.  
 dosage forms. The first method is based on the reaction of the above  
 cited drugs with bromocresol purple dye to form ion-pair complex  
 extractable with chloroform and subsequently measured  
 spectrophotometrically. Secondly, eosin gives with these drugs ion-pair  
 complex, measurable directly without extraction both spectrophotometrically and  
 spectrofluorimetrically. The last method involves the base-catalyzed  
 condensation of mixed anhydrides of organic acids (citric acid/acetic  
 anhydride) where as the tertiary amino group in the above-cited drugs acts  
 as the basic catalyst. The product of condensation is measured  
 spectrophotometrically. All the reaction conditions for the proposed  
 method were studied.  
 AN 2002:742231 CAPLUS  
 DN 138:193370  
 TI Determination of some histamine H1-receptor antagonists in dosage forms  
 AU Gazy, Azza A.; Mahgoub, Hoda; El-Yazbi, F. A.; El-Sayed, M. A.; Youssef,  
 Rasha M.  
 CS Faculty of Pharmacy, Department of Pharmaceutical Analytical Chemistry,  
 University of Alexandria, Alexandria, 21521, Egypt  
 SO Journal of Pharmaceutical and Biomedical Analysis (2002), 30(3), 859-867  
 CODEN: JPBADA; ISSN: 0731-7085  
 PB Elsevier Science B.V.  
 DT Journal  
 LA English  
 IT 79794-75-5, Loratadine  
 RL: ANT (Analyte); ANST (Analytical study)  
 (determination of some H1 antihistamines by fluorometry and spectroscopy)  
 RN 79794-75-5 CAPLUS  
 CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-

benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)



RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 12 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB Four stability-indicating procedures have been suggested for determination of the

non sedating antihistaminic agent loratadine. Loratadine being an ester undergoes alkaline hydrolysis and the corresponding acid derivative is produced as

a degradation product. Its identity was confirmed using IR and MS. The first procedure is based on determination of loratadine by HPLC with detection at wavelength, 250 nm. Mobile phase is acetonitrile:orthophosphoric acid (35:65) using benzophenone as an internal standard. Sensitivity range is 5.00-50.00 µg/mL. Second determination is a densitometric procedure based on determination of loratadine in the presence of its degradate at λ 246 nm using the mobile phase; methanol:ammonia (10:0.15). Sensitivity range is 1.25-7.50 µg per spot. The third procedure is a spectrophotometric one where a mixture of loratadine and its degradate are resolved by first

derivative

ratio spectra. Sensitivity range is found to be 3.00-22.00 µg/mL, upon carrying out the measurements at wavelengths 236, 262.4 and 293.2 nm. The fourth procedure is based on second derivative spectrophotometry, where D2 measurements are carried out at λ 266 nm. The sensitivity range is 3.00-22.00 µg/mL. The validity of the described procedures was assessed by applying the standard addition technique. Statistical anal. of the results have been carried out revealing high accuracy and good precision. The suggested procedures could be used for determination of loratadine both in pure and dosage forms, as well as in the presence of its degradate.

AN 2002:418184 CAPLUS

DN 137:253110

TI Stability indicating methods for the determination of loratadine in the presence of its degradation product

AU El Ragehy, N. A.; Badawey, A. M.; Khateeb, S. Z.-El

CS Faculty of Pharmacy, Department of Analytical Chemistry, Cairo University, Cairo, 11562, Egypt

SO Journal of Pharmaceutical and Biomedical Analysis (2002), 28(6), 1041-1053  
CODEN: JPBADA; ISSN: 0731-7085

PB Elsevier Science B.V.

DT Journal

LA English


IT 79794-75-5, Loratadine

RL: ANT (Analyte); ANST (Analytical study)

(determination of loratadine in presence of degradation product by HPLC and second derivative spectrophotometry)

RN 79794-75-5 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)




RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 13 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN

AB To be considered for further development, lead structures should display the following properties: (1) simple chemical features, amenable for chemical optimization; (2) membership to an established SAR series; (3) favorable patent situation; and (4) good absorption, distribution, metabolism, and excretion (ADME) properties. There are two distinct categories of leads: those that lack any therapeutic use (i.e., "pure" leads), and those that are marketed drugs themselves but have been altered to yield novel drugs. We have previously analyzed the design of leadlike combinatorial libraries starting from 18 lead and drug pairs of structures (S. J. Teague et al. Angew. Chemical, Int. Ed. Engl. 1999, 38, 3743-3748). Here, we report results based on an extended dataset of 96 lead-drug pairs, of which 62 are lead structures that are not marketed as drugs, and 75 are drugs that are not presumably used as leads. We examined the following properties: MW (mol. weight), CMR (the calculated mol. refractivity), RNG (the number of rings), RTB (the number of rotatable bonds), the number of hydrogen bond donors (HDO) and acceptors (HAC), the calculated logarithm of the n-octanol/water partition (CLogP), the calculated logarithm of the distribution coefficient at pH 7.4 (LogD74), the Daylight-fingerprint druglike score (DFPS), and the property and pharmacophore features score (PPFS). The following differences were observed between the medians of drugs and leads:  $\Delta MW = 69$ ;  $\Delta CMR = 1.8$ ;  $\Delta RNG = \Delta HAC = 1$ ;  $\Delta RTB = 2$ ;  $\Delta CLogP = 0.43$ ;  $\Delta LogD74 = 0.97$ ;  $\Delta HDO = 0$ ;  $\Delta DFPS = 0.15$ ;  $\Delta PPFS = 0.12$ . Lead structures exhibit, on the average, less mol. complexity (less MW, less number of rings and rotatable bonds), are less hydrophobic (lower CLogP and LogD74), and less druglike (lower druglike scores). These findings indicate that the process of optimizing a lead into a drug results in more complex structures. This information should be used in the design of novel combinatorial libraries

that are aimed at lead discovery.

AN 2001:527755 CAPLUS  
DN 135:266637  
TI Is There a Difference between Leads and Drugs? A Historical Perspective  
AU Oprea, Tudor I.; Davis, Andrew M.; Teague, Simon J.; Leeson, Paul D.  
CS AstraZeneca R&D Molndal EST Lead Informatics, Moelndal, S 431 83, Swed.  
SO Journal of Chemical Information and Computer Sciences (2001), 41(5),  
1308-1315  
CODEN: JCISD8; ISSN: 0095-2338  
PB American Chemical Society  
DT Journal  
LA English  
IT 38092-89-6  
RL: PRP (Properties)  
(drug design and structure-activity relationship between leads and  
leadlike drugs)  
RN 38092-89-6 CAPLUS  
CN 5H-Benzo[5,6]cyclohepta[1,2-b]pyridine, 8-chloro-6,11-dihydro-11-(1-methyl-  
4-piperidinylidene)- (9CI) (CA INDEX NAME)



RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 14 OF 15 CAPLUS COPYRIGHT 2006 ACS on STN  
AB Introduction of bromine at the 10-position of 3-bromo-8-chlorobenzocycloheptapyridine analogs results in formation of racemic atropisomeric compds. that are easily separable at room temperature on a ChiralPak AD column providing pure atropisomers. Evaluation of the FPT (farnesyl protein transferase) activity of these atropisomers revealed that dextrorotatory compds. were more potent in the farnesyl protein transferase enzyme and cellular assay than their levorotatory isomers. The dextrorotatory were found to inhibit farnesyl protein transferase processing in COS cells at low micro molar range. They were also found to have excellent cellular antitumor activity. Evaluation of the dextrorotatory isomers in DLD-tumor model in nude mice revealed that they were efficacious, inhibiting tumor growth by 55 and 63% at 50 mpk, resp.

AN 1999:383510 CAPLUS  
DN 131:157697  
TI Atropisomeric trihalobenzocycloheptapyridine analogues provide stereoselective FPT inhibitors with antitumor activity  
AU Njoroge, F. George; Vibulbhan, Bancha; Bishop, W. Robert; Kirschmeier, Paul; Bryant, Mathew S.; Nomeir, Amin A.; Liu, Ming; Doll, Ronald J.; Girijavallabhan, Viyyoor M.; Ganguly, Ashit K.  
CS Departments of Chemistry and Tumor Biology, Schering-Plough Research



at the same doses. Dual PAF and histamine antagonist SCH-37370 exhibited mild anti-H1 activity, the maximum effect being 27% at 10 mg/kg. Pure PAF antagonists WEB-2086 and SR-27417 showed no effect against histamine-induced wheals. Only rupatadine, SR-27417A, SCH-37370, and WEB-2086 showed PAF antagonist activity, whereas pure antihistamines were inactive. The most potent PAF antagonist was SR-27417A, with a maximum effect of 56% and 80% at 1 and 10 mg/kg, resp. Rupatadine and WEB-2086 antagonized PAF-induced wheal response, although they showed less maximum effect and shorter duration of action than SR-27417A. SCH-37370 exhibited only slight PAF antagonist activity at 10 mg/kg. Overall, the histamine- and PAF-induced wheal model in dogs proved useful for independent evaluation of histamine and PAF antagonist properties of the tested compds., as pure antagonists blocked the effect of only one of the mediators. Rupatadine was the only of the seven compds. studied that showed potent dual activity against PAF and histamine.

AN 1997:112567 CAPLUS

DN 126:207287

TI Dual effect of a new compound, rupatadine, on edema induced by platelet-activating factor and histamine in dogs: comparison with antihistamines and PAF antagonists

AU Queralt, Mireia; Merlos, Manuel; Giral, Marta; Puigdemont, Anna

CS Departamento de Farmacologia, Facultad de Veterinaria, Universidad Autonoma de Barcelona, Barcelona, Spain

SO Drug Development Research (1996), 39(1), 12-18

CODEN: DDREDK; ISSN: 0272-4391

PB Wiley-Liss

DT Journal

LA English


IT 79794-75-5, Loratadine

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(dual effect of a new compound, rupatadine, on edema induced by platelet-activating factor and histamine in dogs and comparison with antihistamines and PAF antagonists)

RN 79794-75-5 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-, ethyl ester (9CI) (CA INDEX NAME)



=> FIL STNGUIDE

10510619

18/12/2006

Page 23

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	90.94	258.09
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-12.00	-12.00

FILE 'STNGUIDE' ENTERED AT 11:36:04 ON 18 DEC 2006  
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
LAST RELOADED: Dec 8, 2006 (20061208/UP).

10510619